



Urine Collection Kit Manual

The All-in-One
Diagnostic Test
for Infectious Diseases for
All Animals



Catalog No. MI-101
Storage: Room Temperature

MiDOG

Animal Diagnostics

Customer Support:

(833) 456-4364 or (833) iLMiDOG
info@midogtest.com | www.midogtest.com

Orders:

orders@midogtest.com | www.midogtest.com/orders


The information contained in this MiDOG® report is intended only to be factor for use in a diagnosis and treatment regime for the canine patient. As with any diagnosis or treatment regime, you should use clinical discretion with each patient based on a complete evaluation of the patient, including history, physical presentation and complete laboratory data, including confirmatory tests. All test results should be evaluated in the context of the patient's individual clinical presentation. The information in the MiDOG® report has not been evaluated by the U.S. Food and Drug Administration.

CAUTION: Urine Conditioning Buffer™ preservation reagent is provided in the urine collection tube. DO NOT drink, touch, or remove the preservation reagent from the swab collection tube. The preservation reagent is an irritant to the skin and eyes. Do not ingest preservation reagent; avoid contact with skin and eyes; wash with water for 20 minutes if exposed to skin or eyes. Do NOT induce vomiting if the preservation reagent is ingested. Obtain appropriate medical attention in the event of any exposure. Keep out of reach of children.

Clear Plastic Bag



Sample Information Card



Kit Components

Urine Collection Tube



Return Envelope with Prepaid Label




Urine Collection Kit – Instruction Manual



1. Label urine collection tube with the names of the veterinarian and the patient.

Sampling Examples:

Urine collection via cystocentesis



2A. Collect the urine from your patient using cystocentesis.

Urine collection via free-catch



3A. Use a sterile container to catch the urine from the patient.



2B. Transfer collected urine from the syringe into the MiDOG® urine collection tube.



3B. Pour the collected urine into the provided MiDOG® urine collection tube.

Use **ONE** collection tube for each patient.

Pease provide as much urine as possible using the MiDOG® urine collection tube.
IDEALLY PROVIDE AT LEAST 2 ML OF URINE.

Continue (Steps 3-6):




3. Screw the cap securely onto the urine collection tube. Ensure the tube is sealed tightly to prevent leakage.



4. Shake the urine collection tube 10 times to mix the urine with the solution.



5. (A) Place collection tube into the plastic bag. (B) Place the clear plastic bag into the prepaid envelope.



6. (A) Fill out Sample Information Sheet and add to prepaid envelope, or log into your account at www.midogtest.com to provide sample information. (B) Drop the return envelope off at your FedEx location, or schedule a FedEx pickup, within 1 month of sample collection.

After the sample is received, your MiDOG® Report will be sent to the email address you provided within 2 to 5 business days. The report will also be accessible through your MiDOG portal.