



# All-in-One Microbial Test

<b>Patient Name</b>	NA	<b>Health Status</b>	NA	<b>Sample ID</b>	MI1900611
<b>Owner's Name</b>	NA	<b>Ordered By</b>	NA	<b>Sample Type</b>	Left Ear
<b>Gender</b>	Female, Spayed	<b>Email</b>	NA	<b>Received Date</b>	5/7/20
<b>Breed</b>	Bloodhound	<b>Hospital</b>	NA	<b>Report Date</b>	8/26/2020
<b>Age</b>	3 years 5months	<b>Location</b>	NA		
<b>Species</b>	Canine	<b>Account Number</b>	NA		

## Potential Clinically Relevant Microbes Detected:

Listed are those bacteria and fungi detected in the specimen that are of potential clinical relevance. Results from this report should be considered together with clinical data gathered by the veterinarian (physical examination, medical history, cytology, etc.) as the microbes detected may or may not be the cause of the clinical condition. For a comprehensive list of all microorganisms detected in this specimen see page 3 of this report. The purpose of Significance is to highlight those species that are above the expected range for the average clinically healthy dog.

### 1. Bacteria

Species Detected	Relative Abundance (%)	Cells per sample	Reference Range	Significance
<a href="#">Pseudomonas aeruginosa</a>	50.08	130000000	0-249	● High
<a href="#">Streptococcus canis</a> [7]	39.16	100000000	0-1329	● High
<a href="#">Staphylococcus pseudintermedius</a> [2]	2.61	7100000	0-60627	● High
<a href="#">Proteus mirabilis</a> [4]	1.22	3300000	0-0	● High

### 2. Fungi


Species Detected	Relative Abundance (%)	Cells per sample	Reference Range	Significance
<a href="#">Malassezia pachydermatis</a> [8]	95.64	18000	0-2887	● High

#### Abbreviation Key:

- **Normal.** Species detected within the reference range of clinically healthy dogs.
- **Intermediate.** Species detected outside the reference range of clinically healthy dogs.
- **High.** Species detected significantly higher than the reference range of clinically healthy dogs.

## Microbial Overview:

Charts below depict *Bacteria vs Fungi*: an overview of the microbiome, *Bacteria*: the relative abundance of all clinically relevant species detected relative to the rest of the microbiome for bacteria, and *Fungi*: the relative abundance of all clinically relevant species detected relative to the rest of the microbiome for fungi. Each color represents a different species. The larger the size of the colored segment, the more abundant that specific species is in the specimen. The purpose of these graphs is to highlight if any clinically relevant organism is overgrown in the sample.





Patient Name NA  
Owner's Name NA

Ordered By NA  
Account Number NA

## Antibiotic Resistance Panel for Detected Clinically Relevant Microbes

The sample was screened for the presence of antibiotic resistance genes and intrinsic resistances of clinically relevant microorganisms. We recommend the cautious use of any antibiotic drug and to follow the guidelines for antimicrobial stewardship in veterinary practice.

Antibiotics	<i>Pseudomonas aeruginosa</i> (50.08%)	<i>Streptococcus canis</i> (39.16%)	<i>Staphylococcus pseudintermedius</i> (2.61%)	<i>Proteus mirabilis</i> (1.22%)	Suggested Dose	Drug Delivery
Ampicillin	IR	ND	ND	ND		
Amoxicillin	IR	ND	ND	ND		
Oxacillin	ND	ND	ND	ND	--	--
Benzylpenicillin	ND	ND	ND	ND	--	--
Cefadroxil	IR	ND	ND	ND		
Cefazolin	IR	ND	ND	ND		
Cefpodoxime	IR	ND	ND	ND		
Ceftiofur	IR	ND	ND	ND		
Cephalexin	ND	ND	ND	ND	22 mg/kg, q 12 hours	PO
Cefovecin	IR	ND	ND	ND		
Cefoxitin	IR	ND	ND	ND		
Enrofloxacin	ND	ND	ND	ND	5 mg/kg, q 24hrs	PO
Marbofloxacin	ND	ND	ND	ND	2.75 mg/kg, q 24hrs	PO
Ciprofloxacin	ND	ND	ND	ND	--	--
Pradofloxacin	ND	ND	ND	ND	--	--
Clindamycin	IR	ND	ND	IR		
Lincomycin	ND	ND	ND	ND	--	--
Erythromycin	ND	ND	ND	ND	10 mg/kg q 12 hrs	PO
Azithromycin	ND	ND	ND	ND	5 mg/kg q 12 hrs	PO
Sulfonamide	ND	ND	ND	ND	30 mg/kg, q 12hrs	PO
Doxycycline	IR	ND	ND	IR		
Tetracycline	IR	ND	ND	IR		
Minocycline	ND	ND	ND	ND	--	--
Neomycin	IR	IR	ND	ND		
Amikacin	ND	IR	ND	ND		
Gentamicin	ND	IR	ND	ND		
Clavamox	IR	ND	ND	ND		
Timentin	ND	ND	ND	ND	Topical use	Topical use
Ceftazadime	ND	ND	ND	ND	--	--
Florfenicol	ND	ND	ND	ND	--	--
Chloramphenicol	IR	ND	ND	ND		
Imipenem	ND	ND	ND	ND	--	--
Mupirocin	ND	ND	ND	ND	Topical use	Topical use
Rifampicin	ND	ND	ND	ND	--	--

This table lists antibiotic sensitivities/resistances for the indicated bacteria based on detection of specific antibiotic resistance genes and naturally occurring, or intrinsic, resistance to specific antibiotics previously identified for that organism. To receive a list of the antibiotic resistance genes detected as well as intrinsic resistances for additional organisms not listed here, please contact MiDOG® customer support.

### Abbreviation Key:

**ND = Not Detected.** According to our test the organism is most likely sensitive to this antibiotic. No antibiotic resistance gene or intrinsic resistance was detected.

**R = Resistance.** An Antibiotic resistance gene for the indicated antibiotic was detected and the organism is likely resistant to the suggested dose.

**IR = Intrinsic Resistance.** No resistance genes are detected or available for testing, would not recommend this antibiotic due to likely resistant to the indicated antibiotic based on intrinsic factors inferred from microbiological and clinical literature.

**PO = Oral.** By Mouth.

**IV = Intravenous Injection.** Injections involving direct injection into the vein.

**SC = Subcutaneous Injection.** The medication delivered to the tissues between the skin and the muscle.

## Supplemental Data on Microbial Composition

### Bacterial Analysis

Charts below depict the relative abundance of all detected bacterial species. Each color represents a different bacterial species. The larger the size of the colored segment, the more abundant that specific species is in the specimen.



The table below lists all bacterial species detected in the specimen within the limit of detection. The absolute and relative abundances of each species is shown.

Species Detected	Relative Abundance (%)	Cells per sample	Reference Range	Significance
<a href="#">Pseudomonas aeruginosa</a>	50.08	130000000	0-249	● High
<a href="#">Streptococcus canis</a>	39.16	100000000	0-1329	● High

### Fungal Analysis

Charts below depict the relative abundance of all detected fungal species. Each color represents a different fungal species. The larger the size of the colored segment, the more abundant that specific species is in the specimen.



The table below lists all fungal species detected in the specimen within the limit of detection. The absolute and relative abundances of each species is shown.

Species Detected	Relative Abundance (%)	Cells per sample	Reference Range	Significance
<a href="#">Malassezia pachydermatis</a>	95.64	18000	0-2887	● High

#### Abbreviation Key:

- **Normal.** Species detected within the reference range of clinically healthy dogs.
- **Intermediate.** Species detected outside the reference range of clinically healthy dogs.
- **High.** Species detected significantly higher than the reference range of clinically healthy dogs.



Patient Name NA  
Owner's Name NA

Ordered By NA  
Account Number NA

Page 4 of 4

## References

1. Bygott J. M., Malnick H., Shah J.J., Chattaway M.A., Karas J. A. First clinical case of *Corynebacterium auriscanis* isolated from localized dog bite infection. (2008) *Journal of Medical Microbiology*, 57: 899-900
2. Pierezan, F., Olivry, T., Paps, J. S., Lawhon, S. D., Wu, J., Steiner, J. M., et al. The skin microbiome in allergen-induced canine atopic dermatitis. (2016) *Veterinary Dermatology*, 27(5):332-e82
3. Saridomichelakis M.N., Olivry T. An update on the treatment of canine atopic dermatitis. (2016) *The Veterinary Journal*, 207: 29-37
4. Morrissey I., Moyaert H., de Jong A., El Garch F., Klein U., Ludwig C., Thiry J., Youala, M. Antimicrobial susceptibility monitoring of bacterial pathogens isolated from reiratory tract infections in dogs and cats across Europe: ComPath results. (2016) *Veterinary microbiology*, 191:44-53
5. Morrissey I., Moyaert H., de Jong A., El Garch F., Klein U., Ludwig C., Thiry J., Youala, M. Antimicrobial susceptibility monitoring of bacterial pathogens isolated from reiratory tract infections in dogs and cats across Europe: ComPath results. (2016) *Veterinary microbiology*, 191:44-51
6. Ehrlich G. D., Hu F. Z., Sotereanos N., Sewicke J., Parvizi J., Nara P.L., Arciola, C. R. What role do periodontal pathogens play in osteoarthritis and periprosthetic joint infections of the knee. (2014) *J Appl Biomater Funct Mater* 12(1): 13-20
7. Hitzmann A., Bergmann A., Rohde M., Chhatwal G.S., Fulde M. Identification and characterication of the argenine deiminase system of *Streptococcus canis*. *Vet Microbiol*. 2013 Feb 22;162(1):270-7
8. Meason-Smith, C., Diesel, A., Patterson, A. P., Older, C. E., Mansell, J. M., Suchodolski, J. S., & Rodrigues Hoffmann, A. What is living on your dogs skin? Characterization of the canine cutaneous mycobiota and fungal dysbiosis in canine allergic dermatitis. (2015) *FEMS Microbiology Ecology*, 91(12):112
9. Rodrigues Hoffmann A, Patterson AP, Diesel A et al. The skin microbiome in healthy and allergic dogs. (2014) *PLoS One*, 9: e8197

## Method

The MiDOG® All-in-One Microbial Test is a targeted, Next-generation DNA sequencing testing service able to identify molecular signatures unique to the identity and character of a specific microorganism. This test relies on safeguarded preservation and transport of collected samples, thorough extraction of DNA from all microbes present in the specimen, select amplification of microbial DNA followed by Next-generation DNA sequencing using the latest technologies from Illumina (Illumina, Inc., San Diego, CA). Data handling is done via curated microbial databases to accurately align DNA sequences to ensure precise and accurate (species-level) identification of all bacteria and fungi present in the specimen.

## When no Bacterial or Fungal Species are Detected

When no bacterial or fungal species are detected in this test, this result may be due to a very low microbial load and/or low concentration of microbial DNA in the sample provided. In this case, we recommend re-sampling the area of interest and re-submitting specimen for analysis.

## Disclaimer

The information contained in this MiDOG® report is intended only to be factor for use in a diagnosis and treatment regime for the canine patient. As with any diagnosis or treatment regime, you should use clinical discretion with each canine patient based on a complete evaluation of the canine patient, including history, physical presentation and complete laboratory data, including confirmatory tests. All test results should be evaluated in the context of the patients individual clinical presentation. The information in the MiDOG® report has not been evaluated by the FDA.

## Customer Support

Tel: (833)456-4364  
info@midogtest.com  
www.midogtest.com